

# Diagnostic Accuracy of Magnetic Resonance Arthrography and its Correlation with Arthroscopy in the Detection of Lesions Triangular Fibrocartilaginous Complex and Intrinsic Ligaments in the Wrist

Naushaba Malik<sup>1</sup>, Tehmina Sajjad Khan<sup>2</sup>, Shah Bakht Aftab<sup>3</sup>, Minnal Ahmed Malik<sup>4</sup>

Head of Radiology, PESSI Hospital, Islamabad, Consultant Radiologist, Islamabad Diagnostic Center, Sangjani, Rawalpindi  
Post Graduate Trainee, PIMS, Islamabad, House officer, PIMS

## Author's Contribution

<sup>3</sup>Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work and final approval of the version to be published,<sup>2,3</sup>Drafting the work or revising it critically for important intellectual content, <sup>4</sup> literature review

Funding Source: None

Conflict of Interest: None

Received: January 11, 2023

Accepted: March 21, 2023

## Address of Correspondent

Dr Naushaba Malik

Head of Radiology, PESSI Hospital, Islamabad

Doctor.Noshi100@gmail.com

## ABSTRACT

**Objectives:** To assess the diagnostic validity and reliability of magnetic resonance arthrography in identifying the lesion of triangular intrinsic wrist ligaments and fibrocartilaginous complex based on arthroscopy as reference standard and preferred modality of imaging.

**Methodology:** This was a retrospective validation study that was carried out at the Radiology department of PESSI (Punjab Employees Social Security) Hospital, Islamabad, between April 2019 and September 2022. A total of 11 patients with an age range 21-58 years, lesions of the triangular fibrocartilaginous complex (TFCC), and MR arthrography were included. MR arthrography images of the wrist were evaluated to assess the integrity of the intrinsic wrist ligaments, including those of the TFCC, scapholunate, and lunotriquetral ligament. MR arthrography injection was performed into the wrist joint, likely to help visualize the ligaments and other structures within the joint. The dye was injected under fluoroscopic guidance, the size of the needle used for the injection was 20-24G. The data collected from the MR arthrograms and arthroscopy was analyzed using SPSS v 23.

**Results:** The patient's mean age was 40.0±12.31 years. Out of 11 patients, there were 6 (55.5%) males and 5 (45.5%) females. Sensitivity of MR arthrography is 71.4% and specificity that of 75% for detecting TFCC defects, and a 100% sensitivity and 85.7% specificity for diagnosing intrinsic ligament defects. MR arthrography had an overall accuracy of 73% for detecting TFCC, while the overall accuracy for detecting intrinsic ligament defects was 90.9%.

**Conclusion:** MR arthrography was an adequate diagnostic tool for detecting defects of the intrinsic ligaments and the TFCC of the wrist when compared to diagnostic arthroscopy.

**Keywords:** Arthrography, Arthroscopy, Fibrocartilaginous, Ligaments, Magnetic resonance, Wrist joint.

Cite this article as: Malik N, Khan TS, Aftab SR, Malik MA. Diagnostic Accuracy of Magnetic Resonance Arthrography and its Correlation with Arthroscopy in the Detection of Lesions Triangular Fibrocartilaginous Complex and Intrinsic Ligaments in the Wrist. *Ann Pak Inst Med Sci.* 2023; 19(1):29-34. doi. 10.48036/apims.v19i1.769

## Introduction

Imaging the wrist can pose various diagnostic complications for radiologists, particularly when it concerns injuries related to small elements including the intrinsic ligaments and the triangular fibrocartilage complex.<sup>1</sup> The wrist joint has complex anatomy and the minute size of its components can cause pathological

conditions to be initially missed easily. Wrist injuries are highly prevalent, and the subsequent impact on functionality should not be moderated.<sup>2</sup> A wrist injury may involve fractures or ligament tears, depending on factors such as bone strength, wrist position during the incident, and the force applied.<sup>3</sup>

Wrist ligament injuries are most frequently observed in bones of first row, known as the proximal carpal row,

while the two joints in the proximal carpal row that are most susceptible to ligament injuries are the lunotriquetral and the scapholunate.<sup>4</sup> The scapholunate ligament, lunotriquetral ligament, and TFCC are the most often implicated in wrist injuries.<sup>5</sup>

Accurate diagnosis or exclusion of injuries to the intrinsic ligaments (lunotriquetral and scapholunate) and TFCC is essential for appropriate clinical management of wrist injuries.<sup>6</sup> However, in most cases, acute injuries are very painful and swollen, making it difficult to perform a proper physical examination and arrive at an accurate diagnosis based on history and examination alone, specifically for small joints ligaments. In these situations, further diagnostic imaging studies are usually needed to help identify the specific injury and determine the most appropriate treatment plan.<sup>7</sup>

Various imaging techniques, including X-rays, fluoroscopy, CT, ultrasound, and MRI are crucial for diagnosing and assessing wrist injuries that are aimed to be treated non-surgically. In some cases, arthroscopy may be necessary to obtain an immediate and accurate diagnosis.<sup>8</sup> When a physical examination or other imaging methods fail to provide a definitive diagnosis, such as in the case of TFCC or ligament injuries, arthroscopy becomes an essential tool.<sup>9</sup>

The ability to identify and evaluate wrist injuries, particularly those related to the TFCC and intrinsic ligaments, has greatly improved due to advancements in imaging technology over the past two decades.<sup>10</sup> CT, US, MRI, and magnetic resonance arthrography are among the most commonly used imaging techniques for assessing wrist injuries, including those involving the TFCC and intrinsic ligaments.<sup>11</sup>

The rationale of the study was the clinical presentation of wrist injuries can vary depending on the specific injury and its severity. Advanced imaging modalities such as MR arthrography have been shown to be effective in detecting and diagnosing injuries to the intrinsic carpal ligaments and TFCC. These modalities can help guide appropriate clinical management and avoid unnecessary invasive interventions such as diagnostic arthroscopy. The objectives of the study were to assess the diagnostic validity and reliability of magnetic resonance arthrography in identifying the lesion of triangular fibrocartilaginous complex and intrinsic wrist ligaments based on arthroscopy as reference standard and preferred modality of imaging.

## Methodology

This retrospective validation study was carried out from department of radiology of PESSI (Punjab Employees Social Security) hospital, Islamabad between April 2019 and September 2022. Total 12 patients with age range 21-58 years, both gender, lesions of the triangular fibrocartilaginous complex, intrinsic wrist ligaments and willingness MR arthrography were included in the study. Elderly patients ( $\geq 60$  years of age), and lost follow-up were excluded from the study. Patient notes from hospital records were obtained and performed a wrist arthrogram under the supervision of a radiologist by radiographers in diagnosing and managing wrist injuries.

**Injection technique:** During an MR arthrography procedure, gadolinium contrast material was injected into the joint space with a needle.<sup>12</sup> The procedure was watched with a fluoroscope. Patients report less discomfort when they are given injections using needles with a narrow gauge (20–24G). The fluoroscopic spot radiographs were obtained one after the other, and the MR arthrography was performed with a high-field MR system ranging from 1.5T to 3T with a quadrature wrist coil that was specifically constructed for the patient. The MRI scan utilized a sequence consisting of TR 475-632 and TE 15-34, with the section thickness determined by coronal T1-weighted spin-echo and coronal T1-weighted fat suppression spin-echo. Additionally, the scan incorporated coronal turbo spin-echo MRI and T1-weighted coronal turbo spin-echo MRI, which were repeated three times.

**Arthroscopy technique:** Arthroscopy by expert medical professional was performed to evaluate radiocarpal joints. There was a delay of anything from one month to eighteen months between the MR arthrography and the arthroscopy. Placing the patient on the operating table, administering general anaesthesia to them, and immobilising their hand using a Chinese finger trap and a Velcro wrist strap were the steps used to do the procedure. Patient's hands were scrubbed by iodine solution. Finger traps were disinfected before being hung from a ceiling hook using a rod and chain that were provided. The hand and the wrist were covered with a clean paper drape that had adhesive on it. It was decided to increase the pressure of the high-shoulder tourniquet to 250 mmHg, and a counterweight was attached to the hook on the shoulder strap. Conventional radiocarpal arthroscopy was performed on the injured TFCC and ligaments via the standard 3/4 and 6/R portals.<sup>13</sup> Nonetheless, further midcarpal arthroscopy was performed on some patients to examine the scapholunate

and lunotriquetral ligaments. After that, the portals were sealed, the patient's wrist was wrapped with the strap, and the tourniquet was removed from the patient's arm and to get access to the MR arthrography pictures in the PACS (Picture Archiving and Communication System). We looked at the medical histories of all of the patients and got permission from the hospital ethical committee. Age, gender, medical history (including wrist side and symptoms), MR arthrography results, and arthroscopic findings (surgical notes) were all considered.

**Image analysis:** MR arthrography pictures were used to check for partial or full tears in all of the wrist ligaments, including the TFCC. The TFCC, the scapholunate ligament, and the lunotriquetral ligament were all looked at to see if they had any tears. At the point where the TFCC, SL, and LT meet, a contrast leak showed a tear. The number of leaking's showed how bad the defect was (tear). A partial-thickness tear only went through part of the palmar, medial, or dorsal area, while a full-thickness tear went all the way through the area, or the area was lacking.

**Statistical analysis:** Using SPSS v 23, data from MR arthrograms and arthroscopy outcomes were combined. Quantitative data were analyzed by the mean and standard deviation, whereas qualitative data were analyzed by frequencies and percentages. MR arthrograph findings (for TFCC tears and intrinsic ligaments tears) were compared to arthroscopy findings (as documented in each patient's surgical record), taking arthroscopy as the gold standard. Estimates were produced on the specificity, sensitivity and accuracy of tears. Only when the diseased conditions appeared in both imaging and surgical results were there a true positive result. Every false-negative and false-positive data set were subjected to a false-positive analysis.

## Results

MR arthrograms were performed on a total of 12 patients during the months of April 2019 and September 2022 to get comprehensive information on their medical and surgical conditions. As there was no record kept for one patient, he was excluded from the study. The eleven most recent patients were investigated. There were five female patients, which accounts for 45.5% of the total, and there were six male patients, which accounts for 55.5%. Those that were admitted to the hospital had an average age of 40.0±12.31 years old. From the time of the MR arthrography to the operation, the process typically took 7.8 months. We investigated the efficacy of arthroscopic surgery as well as MR arthrography (Table 1).

**Table I: Arthroscopy and MR arthrography for detecting intrinsic ligaments tears and TFCC( N=11)**

	Gender	N	%
	TFCC tear scanned on MR	Male	3
	Female	3	27.3
Ligaments tear on MR	Male	3	27.3
	Female	2	18.2
<b>Total</b>		<b>11</b>	<b>100.0</b>
TFCC tear on arthroscopy	Male	3	27.3
	Female	4	36.3
Ligaments tear on arthroscopy	Male	2	18.2
	Female	2	18.2
<b>Total</b>		<b>11</b>	<b>100.0</b>

Seven individuals who had arthroscopy were discovered to have TFCC tears. Four people had damage to lunotriquetral and scapholunate ligaments. MR arthrography revealed that six individuals had TFCC tears. Five people had damage to the lunotriquetral and scapholunate ligaments. MR arthrography was superior to arthroscopy in locating TFCC tears in 5 of 11 cases (5 true positives and 1 false positive). In one case, arthroscopy detected a TFCC tear that MR arthrography did not (false negative). Arthroscopy did not detect a TFCC rupture in one patient, but MR arthrography detected (false positive).

**Table II: Assessing the diagnostic performance of arthroscopy and MR arthrography in detecting TFCC tears (n=11)**

Test result	Arthroscopy		
	Tear	No tear	Total
<b>MR arthrography</b>	TP = 5	FN = 2	7
	FP = 1	TN = 3	4
	Total = 6	5	11

In five of the seven patients, MR arthrography proved to be a significantly more effective method for detecting TFCC tears compared to arthroscopy. MR arthrography can identify TFCC tear with a sensitivity of 71.4 percent. Meanwhile, 75% is specificity (true negative) rate of MR arthrography. In two patients, MR arthrography could not detect a TFCC tear, with a false positive result, while arthroscopy successfully identified the tear leading to false-negative outcomes for MR arthrography). In one case, MR arthrography detected a tear that arthroscopy had failed to observe, resulting in a false-positive outcome. The research investigated sensitivity, specificity and positive predictive value (PPV), negative predictive value (NPV), and accuracy for detecting TFCC and intrinsic ligament tears (Table III & V).

PPV:  $TP/(TP+FP) \times 100$   
 NPV:  $TN/(FN+TN) \times 100$   
 Accuracy:  $(TP+TN)/(TP+TN+FP+FN) \times 100$   
 Sensitivity:  $TP/(TP+FN) \times 100$   
 Specificity:  $TN/(FP+TN) \times 100$

**Table III: MR arthrography in detecting TFCC tear (n=11)**

Accuracy	72.7%
Negative predictive value	60.0%
Positive predictive value	83.3%
Sensitivity	71.4%
Specificity	75.0%

**Table IV: Comparison of A comparison of the diagnostic performance of MR arthrography and arthroscopy in detecting tears of intrinsic ligaments (n=11)**

Test result	Arthroscopy			
	Tear	No tear	Total	
MR arthrography	Tear	TP: 4	FP: 1	5
	No Tear	FN: 0	TN: 6	6
	Total	4	7	11

**Table V: MR arthrography in detecting intrinsic ligaments tear (n=11)**

Accuracy	90.9%
NPV	100.0%
PPV	80.0%
Sensitivity	100.0%
Specificity	85.7%

The diagnostic accuracy of MR arthrography for TFCC tear detection is 73%. In all four patients, MR arthrography outperformed arthroscopy in identifying tears within the joint. MR arthrography provides a 100% positive predictive value in detecting intrinsic ligament abnormalities (sensitivity). For MR arthrography, 85.7% of negative results were validated. In one case, MR arthrography discovered a tear that arthroscopy had failed to detect (false positive). MR arthrography was successful in identifying a torn intrinsic ligament 91% of the time.

## Discussion

It was seen in 11 MR arthrograms correlated with arthroscopic findings to confirm the TFCC and intrinsic ligaments tears in this study. MR arthrography detects TFCC abnormalities with a sensitivity of 71.4%, and a specificity of 75%. It had a sensitivity of 100 percent and a specificity of 85.5 percent for identifying intrinsic ligament anomalies. 73% accuracy of TFCC, whereas 91% accuracy of ligaments was measured in this study. The average sensitivity of MR arthrography for TFCC tears was 71.4%, which was good for intrinsic tears. 91% tears in the intrinsic ligaments are accurately diagnosed. The typical TFCC tear percentage is 73%. Prior to MR arthrography, arthroscopy was performed on each

participant in this study. In terms of validating the diagnosis and evaluating the reliability and accuracy of this imaging technique, arthroscopy has produced either negative or positive results. On the other hand, the diagnostic accuracy and sensitivity MRA suggest that it is an invasive technique. In some circumstances, diagnostic arthroscopy may not be necessary, although large-scale studies are required to evaluate whether MRA is associated with greater diagnostic sensitivity for TFCC tears. Magnetic resonance imaging (MRI) is believed to have revolutionised the identification and treatment of wrist conditions. Despite this, early research failed to demonstrate that MR arthrography was considerably superior to conventional MRI of the wrist, with sensitivity and specificity of 50% and 50%, respectively.<sup>14</sup> It has been shown that MR arthrography may identify scapholunate ligament lesions 86% of the time. By injecting contrast material into the joint space, MR arthrography of internal structures may be rendered much sharper because the joint capsule expands, intraocular structures can be seen, and tears can be seen because they leak into the joint space above.<sup>15</sup> According to a study by Steinbach et al MR arthrography is the most effective method for evaluating wrist lesions because it combines the benefits of conventional arthrography with magnetic resonance (MR) imaging.<sup>16</sup> Nevertheless, Scheck et al identify and quantify ligamentous anomalies in all regions of the scapholunate ligament, revealing that they correlate well with wrist arthroscopy and have diagnostic and therapeutic applications.<sup>17</sup> Our results are comparable with those of other studies that used MR arthrography to investigate wrist injuries. Several authors compared MR arthrography to different diagnostic techniques to illustrate its sensitivity. According to Brydie et al and Zlatkin et al MRI is a valuable diagnostic technique for identifying injuries to bones, ligaments, and tendons. MRI measurements of the scapholunate and lunotriquetral ligaments were accurate to 90% and 80%, respectively.<sup>18,19</sup> Totterman et al also investigated the sensitivity of MRI for TFCC tear identification.<sup>20</sup> Hobby et al assessed the diagnostic effectiveness of magnetic resonance imaging (MRI). According to the study, MRI had limited reliability in the absence of intrinsic ligament tears, despite displaying high specificity, with only moderate sensitivity. When compared to arthroscopy, MRI demonstrated an overall accuracy of 85%, sensitivity of 70%, and a specificity of 90%.<sup>21</sup> In contrast, Scheck et al showed that MR arthrography had a high level of specificity and sensitivity, ranging from 90% to 96% and 87% to 100%, respectively, for detecting lesions of the intrinsic ligament and TFCC,

with moderate to high sensitivity and support for TFCC connection.<sup>17</sup>

## Conclusion

The results of the study show that magnetic resonance arthrography (MR arthrography) is a better way to find problems in the wrist's intrinsic ligaments and TFCC than diagnostic arthroscopy. When injuries to the scapholunate, lunotriquetral, or TFCC ligaments are suspected, this may be recommended as a risk-free diagnostic method. Even though it will be invasive, and painful. Although it cannot yet replace diagnostic arthroscopy, it may be able to reduce the frequency with which it is required. As MR arthrography enhances picture contrast and diagnostic precision, it should be considered the primary test for those who suspect they have a TFCC or intrinsic ligament injury.

## References

1. Meena T, Roy S. Bone fracture detection using deep supervised learning from radiological images: A paradigm shift. *Diagnostics*. 2022;12(10):2420. <https://doi.org/10.3390/diagnostics12102420>
2. Vassa R, Garg A, Omar IM. Magnetic resonance imaging of the wrist and hand. *Pol J Radiol*. 2020;85(1):461-488. <https://doi.org/10.5114/pjr.2020.99034>
3. Andersson JK. Treatment of scapholunate ligament injury: current concepts. *EFORT Open Rev*. 2017;2(9):382-393. <https://doi.org/10.1302/2058-5241.2.170016>
4. Konopka G, Chim H. Optimal management of scapholunate ligament injuries. *Orthop Res Rev*. 2018;10(2018):41-54. <https://doi.org/10.2147/ORR.S129620>
5. Daunt N, Couzens GB, Cutbush K, Green J, Ross M. Accuracy of magnetic resonance imaging of the wrist for clinically important lesions of the major interosseous ligaments and triangular fibrocartilage complex; correlation with radiocarpal arthroscopy. *Skelet Radiol*. 2021;50(2021):1605-1616. <https://doi.org/10.1007/s00256-020-03701-8>
6. De Santis S, Cozzolino R, Luchetti R, Cazzoletti L. Comparison between MRI and arthroscopy of the wrist for the assessment of posttraumatic lesions of intrinsic ligaments and the triangular fibrocartilage complex. *J Wrist Surg*. 2022;11(01):28-34. <https://doi.org/10.1055/s-0041-1729757>
7. Newton AW, Hawkes DH, Bhalai V. Clinical examination of the wrist. *Orthop Trauma*. 2017;31(4):237-247. <https://doi.org/10.1016/j.mporth.2017.05.009>
8. Know RW. Annual Scientific Meeting Abstracts of the Australasian Musculoskeletal Imaging Group (AMSIG) 2018, Gold Coast, Queensland, Australia. *Skelet Radiol*. 2018;47(2018):1195-1202. <https://doi.org/10.1007/s00256-018-2978-5>
9. Ochman S, Wieskotter B, Langer M, Vieth V, Raschke MJ, Stehling C. High-resolution MRI (3T-MRI) in diagnosis of wrist pain: is diagnostic arthroscopy still necessary?. *Arch Orthop Trauma Surg*. 2017;137(10):1443-1450. <https://doi.org/10.1007/s00402-017-2747-2>
10. Jawed A, Ansari MT, Gupta V. TFCC injuries: How we treat?. *J Clin Orthop Trauma*. 2020;11(4):570-579. <https://doi.org/10.1016/j.jcot.2020.06.001>
11. Cherian BS, Bhat AK, Rajagopal KV, Maddukuri SB, Paul D, Mathai NJ. Comparison of MRI & direct MR arthrography with arthroscopy in diagnosing ligament injuries of wrist. *J Orthop*. 2020;19(5):203-207. <https://doi.org/10.1016/j.jor.2019.11.014>
12. Boer BC, Vestering M, Van Raak SM, van Kooten EO, Huis In't Veld R, Vochteloo AJ. MR arthrography is slightly more accurate than conventional MRI in detecting TFCC lesions of the wrist. *Eur J Orthop Surg Traumatol*. 2018;28(12):1549-1553. <https://doi.org/10.1007/s00590-018-2215-x>
13. Omar NN, Mahmoud MK, Saleh WR, Almallah HG, Qenawy OK, Mourad AF, et al. MR arthrography versus conventional MRI and diagnostic arthroscopy in patients with chronic wrist pain. *Eur J Radiol Open*. 2019;6(1):265-274. <https://doi.org/10.1016/j.eiro.2019.06.003>
14. Kamal HA, Abdelsattar MH, El-Liethy NE. Evaluation of lesions of the internal ligaments of the wrist; conventional magnetic resonance imaging versus MR arthrography (MRA). *Egypt J Radiol Nucl Med*. 2014;45(3):771-778. <https://doi.org/10.1016/j.ejnm.2014.06.005>
15. Lee RK, Ng AW, Tong CS, Griffith JF, Tse WL, Wong C, et al. Intrinsic ligament and triangular fibrocartilage complex tears of the wrist: comparison of MDCT arthrography, conventional 3-T MRI, and MR arthrography. *Skelet Radiol*. 2013;42(9):1277-1285. <https://doi.org/10.1007/s00256-013-1666-8>
16. Steinbach LS, Palmer WE, Schweitzer ME. Special focus session: MR arthrography. *Radiographics*. 2002;22(5):1223-1246. <https://doi.org/10.1148/radiographics.22.5.g02se301223>
17. Scheck RJ, Romagnolo A, Hierner R, Pfluger T, Wilhelm K, Hahn K. The carpal ligaments in MR arthrography of the wrist: correlation with standard MRI and wrist arthroscopy. *J Magn Reson Imaging*. 1999;9(3):468-474.

- [https://doi.org/10.1002/\(SICI\)1522-2586\(199903\)9:3<468:AID-JMRI16>3.0.CO;2-T](https://doi.org/10.1002/(SICI)1522-2586(199903)9:3<468:AID-JMRI16>3.0.CO;2-T)
18. Brydie A, Raby N. Early MRI in the management of clinical scaphoid fracture. *Br J Radiol.* 2003;76(905):296-300.  
<https://doi.org/10.1259/bjr/19790905>
  19. Zlatkin MB, Chao PC, Osterman AL, Schnall MD, Dalinka MK, Kressel HY. Chronic wrist pain: evaluation with high-resolution MR imaging. *Radiology.* 1989;173(3):723-729.  
<https://doi.org/10.1148/radiology.173.3.2813777>
  20. Totterman SM, Miller RJ, McCance SE, Meyers SP. Lesions of the triangular fibrocartilage complex: MR findings with a three-dimensional gradient-recalled-echo sequence. *Radiology.* 1996;199(1):227-232.  
<https://doi.org/10.1148/radiology.199.1.8633149>
  21. Hobby JL, Tom BD, Bearcroft PW, Dixon AK. Magnetic resonance imaging of the wrist: diagnostic performance statistics. *Clin Radiol.* 2001;56(1):50-57.  
<https://doi.org/10.1053/crad.2000.0571>